



**UNIVERSITI PUTRA MALAYSIA**

**PHENOTYPIC AND MOLECULAR CHARACTERIZATIONS OF  
*STREPTOCOCCUS* SPP. ISOLATED FROM BOVINE MAMMARY  
GLANDS**

**MD. FIROZ MIAN**

**FPV 2001 1**

**PHENOTYPIC AND MOLECULAR CHARACTERIZATIONS OF  
*STREPTOCOCCUS SPP.* ISOLATED FROM BOVINE MAMMARY GLANDS**

**By**

**MD. FIROZ MIAN**

**Thesis Submitted in Fulfilment of the Requirement for the Degree of  
Doctor of Philosophy in the Faculty of Veterinary Medicine  
Universiti Putra Malaysia**

**October 2001**



**DEDICATION**

**TO THE MEMORY OF MY BEREAVED FATHER AND TO MY MOTHER  
TO MY WIFE MASUDA AND DAUGHTER FARHIN  
TO MY BROTHER GULAM MUSTAFA**

**AND**

**TO LATE Dr. M. FAZLUR RAHMAN, CHIEF SCIENTIFIC OFFICER, BLRI**

Abstract of the thesis presented to the Senate of the Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**PHENOTYPIC AND MOLECULAR CHARACTERIZATIONS OF  
*STREPTOCOCCUS SPP.* ISOLATED FROM BOVINE MAMMARY GLANDS**  
By

**MD. FIROZ MIAN**

**October 2001**

**Chairman: Abdul Rahim Mutalib, DVM, MS, Ph.D.**

**Faculty: Veterinary Medicine**

Sixty-two streptococcal isolates comprising 20 *Streptococcus agalactiae*, 18 *S. dysgalactiae* and 24 *S. uberis* isolates were recovered from clinical and subclinical cases of bovine mastitis from different dairy herds in the Selangor state in Malaysia. A simple biochemical test scheme formulated on the basis of seven biochemical reactions allowed the identification of *S. agalactiae*, *S. dysgalactiae* and *S. uberis* isolates within 24 hours. *Streptococcus agalactiae* isolates were  $\beta$ -haemolytic, CAMP positive, utilized hippurate, salicin and raffinose; *S. dysgalactiae* isolates were  $\alpha$ -haemolytic and fermented only trehalose and raffinose, while *S. uberis* isolates showed positive reactions to esculin, inulin and mannitol. The API 20 Strep System characterized accurately 100% of *S. agalactiae* and *S. dysgalactiae* isolates, and 96.1% of *S. uberis* isolates although some variable reactions among the isolates within the species were observed. Majority of the isolates were susceptible to most of the antimicrobial agents tested. All *S. agalactiae*, *S. dysgalactiae* and *S. uberis* isolates were susceptible to oxacillin and nitrofurantoin, while 30% of the isolates were resistant to tetracycline. Most of the *S. dysgalactiae* isolates showed resistance

to kanamycin and cephalixin, and *S. agalactiae* isolates to knamycin. *Streptococcus dysgalactiae* isolates showed higher level of resistance compared to *S. agalactiae* and *S. uberis*. Serotyping of the streptococcal isolates using monospecific antisera in agar gel double immunodiffusion revealed that all *S. agalactiae* isolates were typeable and demonstrated the type patterns II (60%), Ia (20%), III (15%) and IV (5%). On the other hand, 17 (94.4%) *S. dysgalactiae* and 22 (91.6%) *S. uberis* isolates were also identified.

The SDS-PAGE and Western-blotting analyses revealed antigenic heterogeneity among the isolates of the bovine *Streptococcus* species examined. Sodium dodecylsulphate polyacrylamide gel electrophoresis could differentiate the three streptococcal species on the basis of their characteristic polypeptide bands. The Western blot analysis also revealed obvious differences in immunogenic proteins between the streptococcal species. Moreover, isolates within each species produced variable protein bands on PAGE analysis and variable immunogenic proteins by Western blotting which let the basis to group them into distinct PAGE and immunoblot fingerprint profiles respectively.

Random amplified polymorphic DNA (RAPD) analysis was evaluated for its capacity to distinguish strains within the species of *S. agalactiae*, *S. dysgalactiae* and *S. uberis* and for epidemiological subtyping. Three single primers were used for each species to generate characteristic RAPD fingerprints. The DNA fingerprint patterns obtained with each primer were distinct and reproducible. The RAPD fingerprints generated could be useful in delineating the strains of *S. agalactiae*, *S. dysgalactiae*

and *S. uberis*. The intraspecies typing efficiency was significantly improved by the parallel use of three primers. The RAPD results showed high level of genetic diversity within strains of the streptococcal species.

The amplification of the DNA encoding 16S rRNA genes by polymerase chain reaction with single set of primers complementary to 16S rRNA gene regions generated characteristic single amplicon that enabled identification and differentiation of the three streptococcal species. The restriction fragment length polymorphism (RFLP) analysis of the amplified 16S rRNA gene regions with the restriction enzymes *MspI* and *RsaI* produced reproducible fingerprint patterns indicating high level of genetic diversity among the isolates of the streptococcal species. Higher heterogeneity was observed within *S. uberis* and *S. dysgalactiae* isolates than the *S. agalactiae* isolates. The discriminatory powers of the two enzymes were to some extent similar.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENCIRIAN FENOTIP DAN MOLEKUL *STREPTOCOCCUS SPP.* YANG  
DIPENCIL DARIPADA KELENJAR MAMA BOVIN**

**Oleh**

**MD. FIROZ MIAN**

**Oktober 2001**

**Pengerusi: Abdul Rahim Mutalib, DVM, MS, Ph.D.**

**Fakulti: Perubatan Veterinar**

Enam puluh dua pencilan streptokokus terdiri daripada 20 *Streptococcus agalactiae*, 18 *S. dysgalactiae* dan 24 *S. uberis* telah diperolehi daripada kes klinikal dan subklinikal mastitis bovin berbagai kelompok lembu tenusu dalam negeri Selangor, Malaysia. Suatu skema ujian biokimia mudah yang dirumus berasaskan tujuh tindak balas, telah membolehkan pengenalpastian pencilan *S. agalactiae*, *S. dysgalactiae*, dan *S. uberis* dalam tempoh 24 jam. Pencilan *Streptococcus agalactiae* ialah  $\alpha$ -hemolisis, positif pada ujian CAMP, pengguna hipurat, salisin dan rafinosa; pencilan *S. dysgalactiae* ialah  $\alpha$ -hemolisis dan hanya menapai trehalosa dan rafinosa, sambil *S. uberis* menunjukkan tindak balas positif terhadap eskulin, inulin, dan manitol. API 20 Strep System mencirikan dengan tepat 100% daripada pencilan *S. agalactiae* dan *S. dysgalactiae*, 96.1% daripada pencilan *S. uberi*, walaupun terdapat pelbagai tindak balas di kalangan pencilan dalam sesuatu spesies. Sebahagian besar daripada pencilan ini rentan terhadap kebanyakan agen antimikrob yang diuji.

Kesemua pencilan *S. agalactiae*, *S. dysgalactiae* dan *S. uberis* adalah rentan terhadap oksasilin dan nitrofurantoin, sambil 30% daripada pencilan tahan tetrasiklin. Kebanyakan daripada pencilan *S. dysgalactiae* menunjukkan ketahanan terhadap kanamisin dan sefalekssin, dan pencilan *S. agalactiae* pula terhadap kanamisin. Pencilan *Streptococcus dysgalactiae* menunjukkan ketahanan lebih tinggi berbanding *S. agalactiae* atau *S. uberis*. Penserotipan pencilan streptokokus mengguna antiserum monokhusus dalam gel agar-agar pengimunoresapan dedua menunjukkan pencilan *S. agalactiae* boleh ditip dan menyatakan pola tip II (60%), III (20%), dan IV (5%). Disebaliknya, 17 (94,4%) pencilan *S. dysgalactiae* dan 22 (91.6%) pencilan *S. uberis* telah dikenal pasti.

Analisis SDS-PAGE dan pensapan Western menunjukkan keheterogenan antigen di kalangan pencilan spesies *Streptococcus bovin* yang dikaji. Elektroforesis gel poliakrilamida natrium dodesilsulfat boleh membeza tiga spesies streptokokus berasaskan jalur polipeptida cirian. Analisis sap Western juga menunjukkan kelainan nyata dalam protein imunogen di antara spesies streptokokus. Tambahan pula pencilan di dalam setiap spesies menghasilkan jalur protein pelbagai pada analisis PAGE dan protein imunogen pelbagai pada pensapan Western, yang mengesahkan asas untuk mengumpulkan pencilan ini masing-masing kepada profil sidikjari PAGE dan imunosap.

Analisis DNA polimorfik terkuat rawak (RAPD) telah dinilai untuk keupayaannya membeza strain di kalangan spesies *S. agalactiae*, *S. dysgalactiae* dan *S. uberis* dan untuk pengsubtipan epidemiologi. Pola sidikjari DNA yang diperolehi daripada setiap primer adalah jelas dan boleh dihasil semula. Sidikjari RAPD yang



dijanakan mungkin berguna dalam membeza di antara strain *S. agalactiae*, *S. dysgalactiae* dan *S. uberis*. Kecekapan pengetipan intraspesies nyata lebih baik dengan penggunaan tiga primer. Hasil RAPD menunjukkan yang aras kepelbagaian genetik di kalangan strain spesies streptokokus adalah tinggi.

Penguatan gen 16S rRNA pengekod DNA melalui tindak balas rangkaian polimerase mengguna satu set primer pelengkap kepada kawasan gen 16S RNA menjanakan amplicon tunggal cirian yang membolehkan untuk pengenapastian dan pembezaan tiga spesies streptokokus dilakukan. Analisis polimorfosime panjang fragmen pengehadan (RFLP) terhadap kawasan gen 16S RNA terkuat dengan mengguna enzim pengehadan *MspI* dan *RsaI* menghasilkan pola sidikjari boleh dihasil semula, menunjukkan yang adanya kepelbagaian genetik aras tinggi di kalangan pencilan spesies streptokokus. Keheterogenan lebih tinggi telah dicerap di kalangan pencilan *S. uberis* dan *S. dysgalactiae* berbanding pencilan *S. agalactiae*. Kuasa penmbezaan untuk dua enzim ini agak sama.

## ACKNOWLEDGEMENTS

All praise to Almighty Allah, the most Merciful and the Benevolent for giving me the will and strength to complete this study.

I would like to express my deepest gratitude and appreciation to Dr. Abdul Rahim Mutalib, Chairman of my supervisory Committee, for his invaluable guidance, constructive comments, advice and valuable suggestions throughout the course of this study.

My utmost appreciation and sincere gratitude to Associate Professor Dr. Abdul Aziz Saharee, Associate Professor Dr. Mohd Azmi Mohd Lila and Dr Nadzri Salim, members of my supervisory committee, for their constructive comments, constant encouragement and invaluable advice towards the completion of this study.

I wish to express my sincere gratefulness to Professor Dr. Abdul Rani Bahaman, Director, Institute of Bioscience and Assoc. Prof. Dr. Mohd Azmi Mohd Lila, Universiti Putra Malaysia for allowing me to use the facilities of their laboratories to conduct my research works.

My special thanks to Professor Dato' Dr. Sheikh Omer Abdul Rahman, Dean of the Faculty of Veterinary Medicine and Professor Dr. Aini Ideris, Dean of Graduate school, Universiti Putra Malaysia for their invaluable advice and moral supports during the study period.

I would like to express my sincere gratitude to Professor Dr. Christoph Lammler, Institute of Milk Hygiene, Justus-Liebig University, Germany, and Dr. James A. Leigh, Compton Lab., UK for providing me with the antisera and reference streptococcal strains used in this study.

I am greatly indebted to the staff members of the faculty, late Mr. Basri Kassim, Mr. Hajaraih Selamat, Mr. Kamaruddin, Mr. Fauzi Che Yousuf, Mr. Krishnan and Mr. Saharuddin for their invaluable technical assistance during the study. I am very grateful to those good friends: Dr. Siti Nor, Mrs Zunita Zakaria, Dr. Siti Khairani Bejo, Dr. Omer Hassan, Mohamad Grissi, Ms Wei Ooi Ling and to those who had contributed directly or indirectly to the completion of this study.

I am most grateful to the Government of the Peoples Republic of Bangladesh and the Director General of Bangladesh Livestock Research Institute (BLRI) for providing me the excellent opportunity with scholarship to accomplish Ph.D.

Last but not least, I wish to express my deepest appreciation to my wife Masuda, whose late night feeding provided the motivation for 'burning the midnight oil' down the home stretch, and to my daughter Farhin for their endurance, constant encouragement and moral supports throughout the course of my study.

I certify that an Examination Committee met on 13<sup>th</sup> October 2001 to conduct the final examination of Md. Firoz Mian on his Doctor of Philosophy thesis entitled "Phenotypic and Molecular Characterizations of *Streptococcus spp.* Isolated from Bovine Mammary Glands" in accordance with Universiti Pertanian Malaysia (Higher degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Abdul Rani Bahaman, DVM, MS, Ph.D.  
Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Chairman)

Abdul Rahim Mutalib, DVM, MS, Ph.D.  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

Mohd Azmi Mohd Lila, DVM, Ph.D.  
Associate Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

Abdul Aziz Saharee, BVSc., MSc., Ph.D.  
Associate Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

Nadzri Salim DVM, MVS, MPVM  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

  
\_\_\_\_\_  
MOHD GHAZALI MOHAYIDIN, Ph.D.  
Professor/ Deputy Dean of Graduate School,  
Universiti Putra Malaysia

Date: 1 NOV 2001

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy.



---

AINI IDERIS, Ph.D.  
Professor,  
Dean of Graduate School,  
Universiti Putra Malaysia

Date: 13 DEC 2001

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or currently submitted for any other degree at Universiti Putra Malaysia or other institutions.

  
MD. FIROZ MIAN

Date: 01-11-2001

# TABLE OF CONTENTS

Page

DEDICATION.....	ii
ABSTRACT.....	iii
ABSTRAK.....	vi
ACKNOWLEDGEMENTS.....	ix
APPROVAL SHEETS.....	xi
DECLARATION FORM.....	xiii
LIST OF TABLES.....	xviii
LIST OF FIGURES.....	xx
LIST OF ABBREVIATIONS.....	xxiv

## CHAPTER

### I

INTRODUCTION.....	1
-------------------	---

### II

LITERATURE REVIEW.....	9
Streptococci.....	9
<i>Streptococcus agalactiae</i> .....	10
<i>Streptococcus dysgalactiae</i> .....	11
<i>Streptococcus uberis</i> .....	11
Bovine Streptococcal Mastitis.....	12
Bacterial Isolation and Identification.....	14
Isolation and Culture Media.....	14
Identification.....	16
Serological Characterization.....	19
Molecular Characterization of Streptococci.....	22
DNA-based Molecular Techniques.....	22
Random Amplified Polymorphic DNA-PCR.....	23
Restriction Enzyme Analysis.....	25
DNA Hybridizations.....	26
Analysis of 16S rRNA gene.....	27
Pulsed-Field Gel Electrophoresis.....	29
Protein-based Molecular Techniques.....	30
Sodium Dodecyl-Sulphate Polyacrylamide Gel Electrophoresis (SDS-PAGE) and Western Blotting.....	30
Antimicrobial Sensitivity.....	33
Virulence Factors and Pathogenicity.....	35
Pathogenesis.....	37
Epidemiology.....	40
Clinical and Pathological Manifestations.....	44
Clinical Manifestations.....	44
Pathological Changes.....	45
Diagnosis.....	46
Clinical Examination and Detection using a strip cup.....	46



Leukocyte and Total Cell Counts.....	47
Indirect Chemical Tests .....	47
Bacteriological Examination.....	47
Serological Diagnosis.....	48
Treatment of Streptococcal Mastitis.....	49
Prevention and Control.....	51
Dry Cow Therapy.....	52
Pre- and Post milking Teat Disinfection.....	52
Antimicrobial therapy.....	53
Vaccination .....	53
Bovine Streptococcal Mastitis in Malaysia.....	56

### III

<b>BIOCHEMICAL AND SEROLOGICAL CHARACTERIZATIONS OF <i>STREPTOCOCCUS SPP.</i> ISOLATED FROM BOVINE MAMMARY GLANDS.....</b>	<b>57</b>
Introduction.....	57
Materials and Methods.....	62
Bacterial Isolation.....	62
Identification of Streptococcal isolates.....	62
API 20 Strep System.....	65
Antibiogram.....	66
Serological Characterization.....	67
Bacterial Cultures.....	67
Preparation of Bacterial Antigens.....	68
Preparation of Monospecific Antisera.....	68
Serotyping Method.....	69
Results.....	70
Isolation and Identification.....	70
The API 20 Strep System.....	73
Antimicrobial Susceptibility Patterns.....	73
Serotyping .....	75
Discussion.....	78

### IV

<b>MOLECULAR CHARACTERIZATION <i>STREPTOCOCCUS SPP.</i> FROM BOVINE MAMMARY GLANDS BY SDS-PAGE AND WESTERN BLOT ANALYSIS.....</b>	<b>86</b>
Introduction.....	86
Materials and Methods .....	90
Bacterial isolates.....	90
SDS-PAGE.....	90
Bacterial Protein Sample Preparation.....	90
Preparation of Hyperimmune Sera in Rabbit.....	91
Gel Components and Preparation.....	91
Electrophoresis Technique.....	92
Gel Staining.....	93



	Western Immunoblotting.....	93
	Protein Transfer to the Nitrocellulose Membrane.....	94
	Immunodetection of the Blotted Proteins.....	95
	Results.....	96
	PAGE-Protein Patterns.....	96
	Western Immunoblotting.....	104
	Discussion.....	112
<b>V</b>	<b>MOLECULAR CHARACTERIZATION OF <i>STREPTOCOCCUS SPP.</i> ISOLATED FROM BOVINE MAMMARY GLANDS BY RANDOM AMPLIFIED POLYMORPHIC DNA-PCR ANALYSIS.....</b>	<b>123</b>
	Introduction.....	123
	Materials and Methods.....	127
	Streptococcal isolates.....	127
	Genomic DNA Preparation.....	127
	Determination of DNA Concentration.....	128
	RAPD-PCR Primers.....	129
	RAPD-PCR Conditions.....	129
	Agarose Gel Electrophoresis .....	130
	Results.....	130
	Discussion.....	148
<b>VI</b>	<b>MOLECULAR CHARACTERIZATION OF <i>STREPTOCOCCUS SPP.</i> FROM BOVINE UDDER INFECTIONS BY ANALYSIS OF 16S rRNA GENE.....</b>	<b>154</b>
	Introduction.....	154
	Materials and Methods.....	158
	Bacterial isolates.....	158
	Isolation of Bacterial DNA.....	158
	PCR Amplification of 16S rRNA gene.....	159
	DNA Primers.....	159
	PCR Conditions.....	159
	Gel Electrophoresis.....	160
	Direct Purification of Amplified 16S rRNA gene.....	160
	Enzymatic Digestions.....	161
	Agarose Gel Electrophoresis.....	162
	16S rRNA gene-RFLP Analysis.....	162
	Results.....	163
	Amplification of 16S rRNA gene by PCR.....	163
	RFLP Analysis of 16S rRNA gene.....	163
	Discussion.....	176
<b>VII</b>	<b>GENERAL DISCUSSION .....</b>	<b>182</b>
	<b>BIBLIOGRAPHY.....</b>	

<b>APPENDICES.....</b>	<b>234</b>
Appendix A: Reagents and Preparations for Bacteriological Techniques.....	235
Appendix B: SDS-PAGE Buffers and Solutions.....	239
Appendix C: Western Blotting Buffers and Reagents.....	245
Appendix D: Reagents and Buffers for DNA Analysis.....	248
 <b>BIO-DATA OF THE AUTHOR.....</b>	 <b>252</b>

## LIST OF TABLES

Table	Page
3 Streptococcal isolates collected from bovine mammary glands from different dairy herds in Selangor, Malaysia.....	71
3.1 Conventional biochemical test scheme for characterization of <i>Streptococcus agalactiae</i> , <i>S. dysgalactiae</i> and <i>S. uberis</i> isolates from bovine intramammary infections .....	73
3.2 Biochemical characteristics (% positive) of <i>Streptococcus agalactiae</i> , <i>S. dysgalactiae</i> and <i>S. uberis</i> isolates utilizing API 20 Strep System.....	74
3.3 Antimicrobial susceptibility patterns of <i>Streptococcus agalactiae</i> , <i>S. dysgalactiae</i> and <i>S. uberis</i> isolated from bovine IMI.....	76
3.4 Serotype distributions of <i>S. agalactiae</i> , <i>S. dysgalactiae</i> and <i>S. uberis</i> isolates recovered from bovine mastitis.....	78
4.1 Number and molecular weights of <i>Streptococcus agalactiae</i> polypeptide bands generated by 12% PAGE stained with coomassie blue.	98
4.2 Number and molecular weights of <i>Streptococcus dysgalactiae</i> polypeptide bands generated by 12% PAGE stained with coomassie blue.....	99
4.3 Number and molecular weights of <i>Streptococcus uberis</i> polypeptide bands generated by 12% PAGE stained with coomassie blue.....	100
5.1 The differences among <i>Streptococcus agalactiae</i> isolates in blotted polypeptide bands immunologically detected by rabbit hyperimmune serum to <i>S. agalactiae</i> .....	106
5.2 The differences among <i>Streptococcus dysgalactiae</i> isolates in blotted polypeptide bands immunologically detected with rabbit hyperimmune serum to <i>S. dysgalactiae</i> .....	107
5.3 The differences among <i>Streptococcus uberis</i> isolates in blotted polypeptide bands immunologically detected with rabbit hyperimmune serum to <i>S. uberis</i> .....	108

6.1	RAPD fingerprint profiles among the <i>S. agalactiae</i> isolates generated with primer OPA-01, OPA-07 and OPA-13.....	133
6.2	RAPD fingerprint profiles among the <i>Streptococcus dysgalactiae</i> isolates generated with primers OPA-01, OPA-07 and OPA-13.....	138
6.3	RAPD fingerprinting profiles of <i>Streptococcus uberis</i> isolates generated with primers OPA-01, OPA-05 and OPA-07.....	143

## LIST OF FIGURES

Figure	Page
3.1 Double immunodiffusion reactions between antisera against serotype II and hot acid extracts of <i>Streptococcus agalactiae</i> isolates (A), and extracts of <i>S. uberis</i> isolates and specific antisera against the reference strain 0140J (B) (two isolates did not react.....	77
4.1 SDS-PAGE protein patterns of <i>Streptococcus agalactiae</i> isolates separated on 12% polyacrylamide gels and stained with coomassie brilliant blue; Lane M: broad range protein molecular standard (kDa).....	101
4.2 SDS-PAGE protein patterns of <i>Streptococcus dysgalactiae</i> isolates separated on 12% polyacrylamide gels and stained with coomassie brilliant blue; Lane M: broad range protein molecular standard (kDa).....	102
4.3 SDS-PAGE protein patterns of <i>Streptococcus uberis</i> isolates separated on 12% polyacrylamide gels and stained with coomassie brilliant blue; Lane M: broad range protein molecular standard (kDa).....	103
5.1 Western immunoblots of whole-cell antigens of <i>Streptococcus agalactiae</i> isolates electrophoresed on 12% SDS-PAGE gels, electrotransferred to nitrocellulose membranes and reacted with rabbit hyperimmune serum to <i>S. agalactiae</i> ; Lane M: broad range protein molecular marker (kDa).....	109
5.2 Western immunoblots of whole-cell antigens of <i>Streptococcus dysgalactiae</i> isolates electrophoresed on 12% SDS-PAGE gels, electrotransferred to nitrocellulose membranes and reacted with rabbit hyperimmune serum to <i>S. dysgalactiae</i> ; Lane M: broad range protein molecular marker (kDa).....	110
5.3 Western immunoblots of whole-cell antigens of <i>Streptococcus uberis</i> isolates electrophoresed on 12% SDS-PAGE gels, electrotransferred to nitrocellulose membranes and reacted with rabbit hyperimmune serum to <i>S. uberis</i> ; Lane M: broad range protein molecular marker (kDa).....	111
6.1 RAPD fingerprinting patterns of <i>Streptococcus agalactiae</i> isolates generated with primer OPA-01 and electrophoresed in 1% agarose gel; Lane M: 1 Kb DNA ladder.....	134

6.2	RAPD fingerprinting patterns of <i>Streptococcus agalactiae</i> isolates generated with primer OPA-07 and electrophoresed in 1% agarose gel; Lane M: 1 Kb DNA ladder.....	135
6.3	RAPD fingerprinting patterns of <i>Streptococcus agalactiae</i> isolates generated with primer OPA-13 and electrophoresed in 1% agarose gel; Lane M: 1 Kb DNA ladder.....	136
6.4	Dendogram of the cluster analysis based on the RAPD fingerprint patterns of <i>Streptococcus agalactiae</i> isolates generated with primers OPA-01, OPA-07 and OPA-13.....	137
6.5	RAPD fingerprinting patterns of <i>Streptococcus dysgalactiae</i> isolates generated with primer OPA-01 and electrophoresed in 1% agarose gel; Lane M: 1 Kb DNA ladder.....	139
6.6	RAPD fingerprinting patterns of <i>Streptococcus dysgalactiae</i> isolates generated with primer OPA-07 and electrophoresed in 1% agarose gel; Lane M: 1 Kb DNA ladder.....	140
6.7	RAPD fingerprinting patterns of <i>Streptococcus dysgalactiae</i> isolates generated with primer OPA-13 and electrophoresed in 1% agarose gel; Lane M: 1 Kb DNA ladder.....	141
6.8	Dendogram of the cluster analysis based on the RAPD fingerprint patterns of <i>Streptococcus dysgalactiae</i> isolates generated with primers OPA-01, OPA-07 and OPA-13.....	142
6.9	RAPD fingerprinting patterns of <i>Streptococcus uberis</i> isolates generated with primer OPA-01 and electrophoresed in 1% agarose gel; Lane M: 1 Kb DNA ladder.....	144
6.10	RAPD fingerprinting patterns of <i>Streptococcus uberis</i> isolates generated with primer OPA-07 and electrophoresed in 1% agarose gel; Lane M: 1 Kb DNA ladder.....	145
6.11	RAPD fingerprinting patterns of <i>Streptococcus uberis</i> isolates generated with primer OPA-05 and electrophoresed in 1% agarose gel; Lane M: 1 Kb DNA ladder.....	146
6.12	Dendogram of the cluster analysis based on the RAPD fingerprint patterns of <i>Streptococcus uberis</i> isolates generated with primers OPA-01, OPA-07 and OPA-05.....	147

7.1	The fragments of the PCR amplified 16S rRNA genes of <i>Streptococcus agalactiae</i> (A), <i>S. dysgalactiae</i> (B) and <i>S. uberis</i> (C); Lane M: 100 bp plus DNA molecular size marker.....	166
7.2	Schematic representation of PCR amplified 16S rRNA gene-RFLP patterns of <i>Streptococcus agalactiae</i> isolates generated by digestions with <i>MspI</i> and <i>RsaI</i> ; Lane M: 100 bp plus DNA molecular size marker.....	167
7.3	The 16S rRNA gene-RFLP fingerprint profiles of <i>Streptococcus agalactiae</i> isolates produced upon digestion with <i>MspI</i> and electrophoresed in 2% agarose gels; Lane M: 100 bp plus DNA molecular size marker .....	168
7.4	The 16S rRNA gene-RFLP fingerprint profiles of <i>Streptococcus agalactiae</i> isolates produced upon digestion with <i>RsaI</i> and electrophoresed in 2% agarose gels; Lane M: 100 bp plus DNA molecular size marker.....	169
7.5	Schematic representation of PCR amplified 16S rRNA gene-RFLP patterns of <i>Streptococcus dysgalactiae</i> isolates generated by digestions with <i>MspI</i> and <i>RsaI</i> ; Lane M: 100 bp plus DNA molecular size marker.....	170
7.6	The 16S rRNA gene-RFLP fingerprint profiles of <i>Streptococcus dysgalactiae</i> isolates produced upon digestion with <i>MspI</i> and electrophoresed in 2% agarose gels; Lane M: 100 bp plus DNA molecular size marker.....	171
7.7	The 16S rRNA gene-RFLP fingerprint profiles of <i>Streptococcus dysgalactiae</i> isolates produced upon digestion with <i>RsaI</i> and electrophoresed in 2% agarose gels; Lane M: 100 bp plus DNA molecular size marker.....	172
7.8	Schematic representation of PCR amplified 16S rRNA gene-RFLP patterns of <i>Streptococcus uberis</i> isolates generated by digestions with <i>MspI</i> and <i>RsaI</i> ; Lane M: 100 bp plus DNA molecular size marker.....	173
7.9	The 16S rRNA gene-RFLP fingerprint profiles of <i>Streptococcus uberis</i> isolates produced upon digestion with <i>MspI</i> and electrophoresed in 2% agarose gels; Lane M: 100 bp plus DNA molecular size marker.....	174

7.10	The 16S rRNA gene-RFLP fingerprint profiles of <i>Streptococcus uberis</i> isolates produced upon digestion with <i>RsaI</i> and electrophoresed in 2% agarose gels; Lane M: 100 bp plus DNA molecular size marker.....	175
------	---	-----



## LIST OF ABBREVIATIONS

Ab	Antibody
AP-PCR	Arbitrary Primed-Polymerase Chain Reaction
bp	Base pair
BSA	Bovine serum albumin
°C	Degree Celcius
cfu	Colony forming unit
cm	Centimeter
CMT	California Mastitis Test
DNA	Deoxyribonucleic acid
dNTP	Deoxy-nucleotide triphosphate
Dr.	Doctor
e.g.	For example
EDTA	Ethelene Diamine tetra-acetate
ELISA	Enzyme Linked Immunosorbent Assay
g	Gram
g/l	Gram per litre
G+C	Guanine + Cytosine
H <sub>2</sub> O <sub>2</sub>	Hydrogen peroxide
h/hrs	Hour/Hours
HS	Hyperimmune Serum
i.e.	That is
i.m.	Intramuscular
IgA	Immunoblobulin A
IgG	Immunoglobulin G
Kbp	Killobase pairs
kDa	Killodalton
M	Molar
Mab	Monoclonal Antibody
M.W.	Molecular Weight
Mg	Milligram
min/mins	Minute/Minutes
ml	Millilitre
mM	Millimolar
nm	Nanometer
O.D.	Optical Density
PAGE	Polyacrylamide Gel Electrophoresis
PBS	Phosphate Buffer Saline
PBST	Phosphate Buffer Saline Tween 20
PCR	Polymerase Chain Reaction
pH	Hydrogen ion concentrartion
PFGE	Pulsed Field Gel Electrophoresis